

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2009**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
White Non-Hispanic	176	41.2	4.4	130	20.7	3.3	0	0.0	0.0	756	23.3	19.0	6,220	23.9	156.3	225	29.0	5.7
Black or African American	29	6.8	11.1	48	7.6	18.4	0	0.0	0.0	823	25.3	316.0	2,414	9.3	926.9	34	4.4	13.1
Hispanic or Latino	171	40.0	9.3	255	40.5	13.9	0	0.0	0.0	851	26.2	46.4	8,653	33.3	471.5	145	18.7	7.9
Asian or Pacific Islander	3	0.7	1.7	7	1.1	4.0	0	0.0	0.0	20	0.6	11.5	262	1.0	150.6	5	0.6	2.9
American Indian or Alaska Native	40	9.4	11.6	43	6.8	12.4	0	0.0	0.0	173	5.3	50.0	2,351	9.0	679.3	109	14.1	31.5
Not Specified	8	1.9	N/A	146	23.2	N/A	0	0.0	NA	627	19.3	N/A	6,102	23.5	N/A	257	33.2	N/A
Total	427	100.0	6.5	629	100.0	9.5	0	0.0	0.0	3,250	100.0	49.3	26,002	100.0	394.2	775	100.0	11.7

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2009.

Note: Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.