

**TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED* YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY
GENDER, RACE/ETHNICITY AND URBAN/RURAL AREAS**, ARIZONA, 2011**

	ARIZONA			Area/gender						Race/ethnicity					
	Total	Male	Female	Total Urban	Urban Male	Urban Female	Total Rural	Rural Male	Rural Female	White Non-Hispanic	Hispanic or Latino	Black or African American	American Indian or Alaska Native	Asian or other Pacific Islander	
Total, all causes	53.8%	61.8%	45.1%	52.7%	60.7%	44.1%	57.7%	65.1%	49.0%	49.6%	67.4%	74.4%	77.7%	66.3%	
Major cardiovascular diseases	42.1%	52.1%	31.4%	40.6%	50.7%	30.2%	47.2%	56.7%	35.9%	38.5%	52.8%	72.1%	66.9%	62.3%	
Diseases of heart	43.3%	53.0%	31.6%	41.9%	51.6%	30.5%	48.1%	57.4%	35.8%	40.1%	54.1%	74.2%	66.5%	62.7%	
Coronary heart disease	45.3%	54.8%	32.6%	43.7%	53.3%	31.3%	51.4%	59.7%	38.6%	42.5%	54.9%	72.6%	63.4%	65.7%	
Malignant neoplasms	66.6%	67.3%	65.8%	66.3%	67.1%	65.4%	67.3%	67.4%	67.1%	64.8%	73.4%	81.5%	73.5%	74.8%	
Lung cancer	68.4%	69.9%	66.5%	68.1%	69.6%	66.2%	69.2%	70.3%	67.7%	68.6%	62.6%	77.6%	61.5%	60.9%	
Prostate cancer	44.4%	44.4%	NA	46.6%	46.7%	NA	35.6%	35.6%	NA	42.9%	52.5%	53.8%	30.0%	66.7%	
Breast cancer	72.8%	66.7%	72.8%	73.4%	33.3%	73.6%	70.1%	100.0%	69.5%	69.2%	91.2%	90.9%	83.3%	92.3%	
Colorectal cancer	63.0%	68.4%	57.1%	62.9%	69.4%	56.3%	62.7%	64.4%	60.5%	59.6%	76.3%	70.0%	85.7%	76.5%	
Malignant melanoma of skin	74.2%	69.6%	83.6%	72.7%	66.3%	84.2%	84.0%	85.7%	75.0%	73.1%	81.8%	0.0%	100.0%	100.0%	
Cerebrovascular diseases	37.0%	46.9%	30.7%	35.5%	44.7%	29.7%	42.1%	52.8%	34.6%	31.8%	50.8%	65.4%	62.2%	63.3%	
Chronic lower respiratory diseases	50.7%	55.3%	46.5%	48.9%	53.2%	45.0%	56.8%	62.5%	51.5%	50.0%	53.3%	68.0%	54.5%	47.1%	
Accidents (unintentional injuries)	75.7%	84.2%	62.5%	73.4%	82.6%	59.4%	82.9%	89.1%	73.2%	69.3%	86.5%	89.9%	95.9%	78.6%	
Motor vehicle accidents	93.2%	94.6%	89.7%	93.2%	94.7%	89.3%	93.1%	94.2%	90.5%	89.8%	95.5%	96.6%	99.3%	90.9%	
Falls	29.6%	43.8%	18.4%	27.3%	41.3%	16.3%	40.9%	56.3%	29.9%	25.5%	46.6%	40.0%	74.1%	33.3%	
Accidental poisoning	99.4%	99.3%	99.7%	99.3%	99.1%	99.6%	100.0%	100.0%	100.0%	99.3%	99.3%	100.0%	100.0%	100.0%	
Accidental drowning	92.7%	92.7%	92.6%	91.3%	90.9%	92.0%	100.0%	100.0%	100.0%	91.7%	93.8%	83.3%	100.0%	100.0%	
Influenza and pneumonia	39.1%	44.7%	33.6%	38.8%	43.4%	34.6%	40.1%	47.4%	30.7%	34.8%	52.4%	73.3%	43.9%	55.6%	
Alzheimer's disease	13.3%	18.1%	10.9%	13.1%	16.8%	11.2%	14.3%	26.7%	8.0%	12.6%	15.1%	14.8%	40.0%	17.6%	
Diabetes	65.3%	69.8%	59.9%	65.0%	69.5%	59.6%	66.0%	70.4%	60.2%	60.1%	72.1%	78.7%	81.9%	66.7%	
Intentional self-harm (suicide)	93.9%	93.2%	96.3%	94.0%	93.2%	96.8%	93.8%	93.7%	94.4%	92.8%	97.8%	95.8%	100.0%	100.0%	
Chronic liver disease and cirrhosis	92.5%	94.8%	88.2%	91.8%	94.5%	86.8%	94.6%	95.8%	92.5%	91.0%	93.5%	100.0%	96.5%	100.0%	
Nephritis, nephrotic syndrome, and nephrosis	42.7%	46.6%	38.6%	43.0%	47.9%	37.2%	41.7%	42.4%	41.2%	35.4%	56.9%	75.0%	64.0%	55.6%	
Septicemia	58.8%	62.2%	55.2%	61.6%	63.2%	59.6%	52.4%	59.0%	46.7%	52.7%	66.0%	90.9%	72.4%	75.0%	
Assault (homicide)	97.6%	98.3%	95.1%	97.4%	98.4%	93.7%	100.0%	100.0%	100.0%	96.2%	98.6%	100.0%	100.0%	85.7%	
Parkinson's disease	30.9%	34.2%	26.0%	30.2%	33.8%	24.6%	34.0%	35.8%	31.8%	28.8%	48.5%	42.9%	53.3%	0.0%	
Essential (primary) hypertension and hypertensive renal disease	37.2%	48.3%	29.4%	36.3%	46.9%	28.6%	43.4%	60.0%	34.0%	31.1%	48.8%	72.7%	78.6%	60.0%	
Human immunodeficiency virus (HIV) disease	99.0%	98.8%	100.0%	98.8%	98.5%	100.0%	100.0%	100.0%	100.0%	97.9%	100.0%	100.0%	100.0%	100.0%	
INJURY BY FIREARMS	93.3%	93.0%	95.3%	93.4%	93.2%	94.9%	93.0%	92.4%	96.7%	91.1%	98.4%	97.7%	100.0%	100.0%	
DRUG-INDUCED DEATHS	99.1%	99.3%	98.9%	99.1%	99.4%	98.6%	99.4%	98.8%	100.0%	99.4%	98.2%	97.1%	100.0%	100.0%	
ALCOHOL-INDUCED DEATHS	97.2%	97.1%	97.7%	96.9%	96.8%	97.3%	98.0%	97.7%	98.7%	96.3%	98.2%	100.0%	98.9%	100.0%	

* Expected years of life at birth for all U.S. residents.

**Urban = Maricopa, Pima, Pinal, and Yuma counties.