

TABLE 2B-2 (correction)
AGE-ADJUSTED MORTALITY RATES^a BY GENDER AND YEAR FOR SELECTED LEADING CAUSES OF DEATH,
ARIZONA, 2002-2012

		2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
TOTAL, ALL CAUSES												
	Total	796.4	784.0	757.3	772.5	727.3	663.2	659.6	653.2	679.9	699.6	687.2
	Male	964.8	942.8	909.0	925.7	861.6	773.8	777.6	763.9	802.0	817.1	811.0
	Female	658.0	651.5	632.0	644.9	611.8	561.7	554.3	555.1	572.6	595.3	579.2
CARDIOVASCULAR DISEASE												
	Total	265.5	258.4	248.5	245.7	216.4	194.4	194.1	188.8	188.6	201.0	193.3
	Male	321.1	310.3	298.4	298.2	258.3	227.5	235.8	228.1	225.1	239.2	233.6
	Female	221.0	215.8	207.8	203.8	181.0	164.3	158.1	154.9	156.7	168.0	159.1
-----DISEASES OF HEART												
	Total	201.0	197.1	186.7	186.9	166.1	149.5	149.3	145.9	143.3	152.7	145.8
	Male	253.6	246.6	234.2	237.4	207.3	184.0	189.3	183.1	179.8	192.8	184.4
	Female	158.9	157.1	148.5	147.3	131.8	119.4	115.1	114.1	112.2	119.0	113.7
-----CEREBROVASCULAR DISEASE												
	Total	47.0	43.9	43.6	40.6	34.8	30.6	30.3	28.9	30.7	30.6	29.9
	Male	47.3	42.9	42.7	40.0	33.4	27.7	29.6	28.2	28.9	28.0	30.0
	Female	46.7	43.8	43.6	40.5	35.2	32.2	30.5	29.1	31.3	32.3	29.4
MALIGNANT NEOPLASMS												
	Total	168.6	169.4	164.7	161.8	154.7	148.0	145.5	147.2	150.5	151.3	149.8
	Male	209.1	208.3	201.2	197.1	187.0	176.7	173.8	174.9	180.6	179.3	180.3
	Female	139.1	141.3	138.6	136.1	131.0	125.6	123.0	125.8	127.1	129.5	125.9
CHRONIC LOWER RESPIRATORY												
	Total	47.0	45.4	41.9	46.8	44.3	39.2	42.1	40.5	42.6	45.9	44.2
	Male	57.1	53.8	48.1	51.3	50.2	43.5	46.5	44.7	48.1	49.7	50.5
	Female	40.3	40.0	37.7	43.5	40.1	36.0	38.9	37.1	38.6	42.8	39.5
ACCIDENTS (UNINTENTIONAL)												
	Total	45.2	44.8	46.0	51.1	51.2	46.6	44.7	43.1	43.8	45.5	42.4
	Male	62.8	61.6	61.1	68.7	68.5	60.6	59.0	56.6	56.6	58.3	54.0
	Female	28.8	28.8	31.7	34.4	34.3	32.3	30.6	29.8	31.6	32.6	31.0
-----MOTOR VEHICLE ACCIDENTS												
	Total	18.5	18.0	18.1	18.9	19.6	16.2	13.6	11.6	11.1	12.1	11.4
	Male	25.5	24.9	23.9	27.0	26.8	22.2	19.5	16.8	15.9	17.6	16.8
	Female	11.6	11.2	12.5	11.1	12.4	10.1	7.9	6.5	6.4	6.9	6.3
INFLUENZA AND PNEUMONIA^b												
	Total	25.0	23.4	20.1	22.3	18.4	12.9	15.6	14.6	10.9	9.7	9.1
	Male	27.8	29.8	25.7	26.6	22.2	14.9	17.7	15.6	12.8	11.5	7.4
	Female	22.9	18.7	16.3	19.2	15.7	11.5	14.1	13.7	9.3	8.2	5.6

TABLE 2B-2 (continued)
AGE-ADJUSTED MORTALITY RATES^a BY GENDER AND YEAR FOR SELECTED LEADING CAUSES OF DEATH,
ARIZONA, 2002-2012

		2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
ALZHEIMER'S DISEASE^b												
	Total	28.0	32.5	31.3	32.7	33.4	29.6	29.8	29.5	35.1	35.2	29.9
	Male	23.4	26.6	26.4	27.7	27.0	22.8	23.9	24.5	28.5	28.9	41.5
	Female	30.5	35.9	34.2	35.7	47.4	34.1	33.6	32.8	39.3	39.0	52.8
DIABETES												
	Total	22.3	20.3	20.7	20.1	18.9	17.0	17.0	15.7	20.1	24.8	23.5
	Male	25.6	23.6	23.4	24.7	22.5	20.5	20.7	18.2	24.9	30.3	29.2
	Female	19.7	17.5	18.4	16.7	15.9	13.9	13.9	13.5	16.0	20.2	18.6
CHRONIC LIVER DISEASE AND												
	Total	12.1	11.5	11.1	12.7	11.4	11.3	11.7	12.2	12.3	13.4	13.6
	Male	15.6	16.2	15.6	17.7	14.3	15.4	15.3	15.4	16.1	18.0	18.7
	Female	8.9	7.1	6.9	8.1	8.5	7.6	8.4	9.1	8.8	9.2	9.0
NEPHRITIS, NEPHROTIC SYNDROME,												
	Total	11.3	10.1	11.0	10.4	8.8	7.6	7.1	7.7	7.6	6.0	6.2
	Male	12.9	12.0	13.4	12.7	10.5	8.8	8.6	9.2	9.0	7.2	8.9
	Female	10.4	8.8	9.4	8.8	7.4	6.6	5.9	6.5	6.4	5.1	6.7
INTENTIONAL SELF-HARM (SUICIDE)												
	Total	15.9	14.6	14.9	15.4	15.4	15.4	14.8	16.1	16.7	17.2	16.2
	Male	26.4	24.0	24.1	24.9	24.7	24.4	23.0	24.6	27.1	27.4	25.9
	Female	6.0	5.8	6.3	6.5	6.6	6.7	7.0	8.1	6.7	7.5	7.0
ASSAULT (HOMICIDE)												
	Total	8.6	8.3	8.2	8.2	8.5	7.7	7.1	5.5	6.5	6.1	5.9
	Male	13.9	12.8	13.3	13.5	13.5	11.8	11.2	8.0	10.5	9.6	8.6
	Female	3.0	3.6	2.9	2.6	3.1	3.2	2.7	2.9	2.4	2.6	3.1
SEPTICEMIA^b												
	Total	8.3	8.4	8.0	8.4	8.6	8.1	7.9	7.7	7.4	4.7	4.5
	Male	9.6	9.4	9.4	9.7	9.4	8.4	9.4	8.2	10.8	5.4	6.1
	Female	7.2	7.6	7.0	7.5	8.0	7.9	6.7	7.3	5.1	4.0	4.8

Notes: ^a The number of deaths per 100,000 population adjusted to the 2000 U.S. standard. The cause-of-death titles are according to the Tenth Revision of the International Classification of Diseases (ICD-10). Beginning in 2000, the causes of death are classified by ICD-10. The causes of death for 1998-1999 are classified by the Ninth Revision (ICD-9). The rates for 1998-1999 were revised in this edition of the report for consistency with the newly recomputed mortality rates in Table 2B-3. ^b The rates for 1998-1999 are comparability-modified. In order to obtain comparability-modified data, the number of deaths or the death rate for a particular cause of death (such as Alzheimer's disease, Nephritis, Influenza and pneumonia, or Septicemia) classified by ICD-9 is adjusted by multiplying it by the comparability ratio for that cause. Comparability ratios are measures of comparison between ICD-9 and ICD-10. Comparability ratio of 1.0 indicates that the same number of deaths would be assigned to a cause-of-death when ICD-9 or ICD-10 was used. The comparability ratios for causes of death other than the four selected above are within 1.0-1.1 range (see Technical Notes).