

TABLE 2B-7 (correction)
NUMBER OF DEATHS FROM DISEASES OF THE HEART BY CATEGORY, GENDER, AND YEAR, ARIZONA, 2002-2012

		2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
ALL FORMS OF HEART DISEASE	TOTAL	10,551	10,649	10,402	10,779	10,362	10,147	10,299	10,151	9,719	10,424	10,366
	Male	5,642	5,649	5,543	5,789	5,538	5,455	5,701	5,558	5,351	5,748	5,731
	Female	4,909	5,000	4,859	4,990	4,834	4,692	4,598	4,593	4,368	4,675	4,635
Acute rheumatic fever and chronic rheumatic heart disease	TOTAL	46	63	46	55	57	38	57	51	42	69	51
	Male	14	25	18	16	15	15	24	16	15	22	19
	Female	32	38	28	39	42	23	33	35	27	47	32
Hypertensive heart and renal disease^a	TOTAL	31	33	32	39	24	29	48	35	36	49	51
	Male	14	20	9	18	13	15	22	18	16	22	28
	Female	17	13	23	21	11	14	26	17	20	27	23
Ischemic heart disease	TOTAL	7,962	8,037	7,836	8,060	7,628	7,163	7,487	7,372	7,043	7,498	7,294
	Male	4,449	4,402	4,344	4,509	4,202	4,005	4,337	4,233	4,076	4,409	4,279
	Female	3,513	3,635	3,492	3,551	3,426	3,158	3,150	3,139	2,967	3,089	3,015
Hypertensive heart disease	TOTAL	431	440	463	487	506	646	675	698	644	739	761
	Male	197	215	223	226	256	332	348	340	302	348	375
	Female	234	225	240	261	250	314	327	358	342	391	386
Other heart diseases	TOTAL	2,104	2,076	2,025	2,138	2,147	2,271	2,032	1,995	1,954	2,069	2,209
	Male	968	987	949	1,020	1,052	1,088	970	951	942	947	1,030
	Female	1,136	1,089	1,076	1,118	1,095	1,183	1,062	1,044	1,012	1,121	1,179

Notes: Beginning in 2000, the causes of death are classified by ICD-10, replacing the Ninth Revision (ICD-9) used during 1979-1999; ^a The comparability ratio for hypertensive heart and renal disease is 1.5, indicating a 50 percent increase in hypertensive heart and renal disease deaths when classified by ICD-10.