

TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED^a YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY
GENDER, RACE/ETHNICITY, AND URBAN/RURAL AREAS^b, ARIZONA, 2012

	ARIZONA			Area/gender						Race/ethnicity				
	Total	Male	Female	Total Urban	Urban Male	Urban Female	Total Rural	Rural Male	Rural Female	White Non-Hispanic	Hispanic or Latino	Black or African American	American Indian or Alaska Native	Asian or Pacific Islander
Total, all causes	55.0%	62.3%	46.7%	53.4%	60.9%	45.4%	58.4%	64.9%	49.9%	50.9%	68.4%	74.4%	78.7%	62.3%
Major cardiovascular diseases	43.7%	53.2%	32.9%	41.4%	50.8%	31.3%	48.4%	57.6%	36.7%	40.5%	55.7%	65.6%	67.6%	57.4%
Diseases of heart	45.0%	54.1%	33.5%	42.4%	51.3%	31.7%	50.8%	59.4%	38.7%	42.3%	55.3%	70.3%	65.8%	55.5%
Coronary heart disease	47.3%	56.3%	34.6%	44.0%	52.9%	32.3%	55.3%	63.6%	41.7%	44.8%	56.6%	70.1%	65.9%	52.5%
Malignant neoplasms	68.4%	69.0%	67.6%	67.7%	68.4%	66.9%	69.6%	70.0%	69.0%	66.5%	74.9%	83.0%	77.9%	79.2%
Lung cancer	70.9%	71.6%	70.2%	70.4%	71.1%	69.7%	73.2%	73.4%	72.9%	70.5%	70.4%	82.3%	66.7%	85.3%
Prostate cancer	47.1%	47.0%	NA	45.2%	45.1%	NA	52.0%	52.0%	NA	45.8%	47.1%	76.2%	66.7%	0.0%
Breast cancer	73.9%	50.0%	74.0%	72.6%	50.0%	72.7%	77.7%	0.0%	77.7%	71.6%	82.4%	87.5%	80.0%	81.3%
Colorectal cancer	66.3%	72.1%	59.1%	67.6%	73.3%	60.8%	61.3%	69.9%	49.4%	63.4%	81.5%	62.5%	88.2%	78.6%
Malignant melanoma of skin	74.8%	73.5%	77.5%	72.9%	73.0%	72.5%	84.6%	76.2%	94.4%	73.9%	91.7%	100.0%	0.0%	100.0%
Cerebrovascular diseases	38.4%	49.2%	30.0%	37.8%	49.0%	29.5%	35.4%	46.8%	24.9%	32.6%	56.6%	57.3%	75.5%	60.0%
Chronic lower respiratory diseases	50.9%	53.1%	48.8%	49.8%	51.5%	48.1%	53.8%	56.4%	50.9%	50.2%	57.5%	75.4%	53.1%	21.4%
Accidents (unintentional injuries)	77.0%	83.6%	66.2%	74.4%	81.4%	63.7%	80.8%	87.4%	70.6%	70.7%	89.2%	91.7%	95.7%	82.5%
Motor vehicle accidents	94.2%	94.6%	93.2%	94.0%	94.6%	92.1%	95.1%	94.3%	96.9%	91.5%	97.2%	96.6%	99.2%	92.9%
Falls	35.1%	42.9%	28.0%	32.8%	39.5%	27.0%	41.4%	53.7%	30.8%	31.5%	48.0%	70.0%	75.0%	50.0%
Accidental poisoning	99.2%	99.4%	98.8%	99.1%	99.2%	98.8%	99.4%	100.0%	98.5%	99.2%	100.0%	100.0%	100.0%	71.4%
Accidental drowning	92.5%	92.3%	92.9%	92.3%	93.6%	88.9%	94.1%	92.9%	100.0%	87.9%	95.5%	100.0%	100.0%	100.0%
Influenza and pneumonia	38.3%	39.8%	37.0%	37.9%	39.2%	36.8%	37.9%	39.3%	36.3%	36.0%	47.3%	58.3%	48.1%	0.0%
Alzheimer's disease	14.7%	18.7%	12.6%	14.5%	17.8%	12.7%	17.0%	27.6%	11.6%	14.0%	22.6%	14.3%	17.6%	6.9%
Diabetes	67.0%	70.9%	61.8%	67.1%	71.5%	61.7%	65.9%	68.6%	61.3%	63.5%	71.8%	72.9%	77.8%	55.9%
Intentional self-harm (suicide)	93.9%	93.0%	97.0%	93.7%	92.8%	96.7%	93.5%	92.5%	98.0%	92.5%	98.4%	100.0%	100.0%	100.0%
Chronic liver disease and cirrhosis	92.4%	92.4%	92.3%	91.4%	91.4%	91.4%	94.1%	94.4%	93.5%	91.8%	91.8%	96.0%	94.7%	100.0%
Nephritis, nephrotic syndrome and nephrosis	42.4%	47.4%	37.2%	40.6%	41.6%	39.7%	44.2%	54.4%	32.4%	36.0%	54.8%	64.3%	57.7%	75.0%
Septicemia	56.6%	63.2%	49.4%	57.9%	65.5%	49.5%	52.1%	54.3%	50.0%	54.6%	60.0%	63.6%	68.8%	42.9%
Assault (homicide)	96.6%	98.0%	92.5%	97.6%	99.5%	91.9%	98.7%	100.0%	96.3%	91.9%	100.0%	100.0%	100.0%	100.0%
Parkinson's disease	27.6%	29.9%	23.6%	27.4%	29.5%	23.7%	27.7%	30.0%	23.5%	26.4%	40.0%	75.0%	33.3%	16.7%
Essential (primary) hypertension and hypertensive renal disease	40.2%	50.6%	31.8%	40.0%	50.9%	31.0%	40.7%	47.9%	35.4%	34.9%	57.1%	50.0%	71.4%	61.5%
Human immunodeficiency virus (HIV) disease	99.0%	100.0%	92.9%	98.9%	100.0%	91.7%	100.0%	100.0%	100.0%	98.3%	100.0%	100.0%	100.0%	100.0%
INJURY BY FIREARMS	93.6%	93.5%	94.6%	94.3%	94.2%	94.8%	93.1%	92.5%	96.6%	91.5%	98.8%	100.0%	100.0%	100.0%
DRUG-INDUCED DEATHS	98.0%	98.8%	96.8%	97.5%	98.6%	95.9%	99.5%	99.1%	100.0%	97.4%	100.0%	100.0%	100.0%	100.0%
ALCOHOL-INDUCED DEATHS	96.2%	95.8%	97.5%	95.7%	95.1%	97.4%	97.4%	97.4%	97.4%	95.8%	95.4%	95.5%	98.1%	100.0%

Notes: ^a Expected years of life at birth for all U.S. residents (78.7 years in 2011); ^b Urban = Maricopa, Pima, Pinal and Yuma counties.