

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, CHLAMYDIA, AND HERPES BY RACE/ETHNICITY, ARIZONA, 2012**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
White Non-Hispanic	148	41.9	3.9	90	21.1	2.4	0	0.0	0.0	1,179	20.1	31.0	5,647	18.5	148.7	503	29.2	13.2
Black or African American	35	9.9	12.2	44	10.3	15.3	0	0.0	0.0	1,091	18.6	380.6	2,535	8.3	884.3	147	8.5	51.3
Hispanic or Latino	125	35.4	6.3	230	53.9	11.6	0	0.0	0.0	1,774	30.3	89.8	10,673	34.9	540.1	385	22.4	19.5
Asian or Pacific Islander	7	2.0	3.2	6	1.4	2.8	0	0.0	0.0	54	0.9	25.1	230	0.8	106.8	14	0.8	6.5
American Indian or Alaska Native	25	7.1	9.1	31	7.3	11.2	0	0.0	0.0	513	8.8	185.7	3,030	9.9	1096.9	129	7.5	46.7
Not Specified	13	3.7	N/A	26	6.1	N/A	0	0.0	N/A	1,245	21.3	N/A	8,456	27.7	N/A	544	31.6	N/A
Total	353	100.0	5.4	427	100.0	6.5	0	0.0	0.0	5,856	100.0	89.4	30,571	100.0	466.5	1,722	100.0	26.3

Notes: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2012; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.