

**TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED^a YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY
GENDER, RACE/ETHNICITY, AND URBAN/RURAL AREAS^b, ARIZONA, 2013**

	ARIZONA			Area/gender						Race/ethnicity				
	Total	Male	Female	Total Urban	Urban Male	Urban Female	Total Rural	Rural Male	Rural Female	White Non-Hispanic	Hispanic or Latino	Black or African American	American Indian or Alaska Native	Asian or Pacific Islander
Total, all causes	54.6%	62.0%	46.2%	53.0%	60.8%	44.6%	57.7%	64.4%	49.7%	50.1%	67.1%	72.8%	76.7%	58.9%
Major cardiovascular diseases	43.2%	53.1%	31.6%	41.4%	51.4%	30.2%	46.4%	56.3%	34.0%	39.5%	53.8%	68.1%	61.2%	52.4%
Diseases of heart	44.1%	53.7%	31.4%	42.2%	51.9%	29.9%	47.4%	57.1%	34.0%	40.5%	55.3%	69.0%	60.2%	51.5%
Coronary heart disease	45.3%	55.3%	31.1%	42.7%	53.0%	28.8%	51.2%	60.0%	37.4%	42.0%	55.2%	67.4%	59.5%	52.6%
Malignant neoplasms	66.8%	67.4%	66.2%	65.9%	66.9%	64.8%	69.0%	67.7%	70.6%	64.8%	74.4%	78.6%	72.4%	66.9%
Lung cancer	68.6%	70.4%	66.6%	67.0%	69.8%	63.9%	73.0%	71.1%	75.2%	68.6%	62.2%	83.1%	55.6%	61.3%
Prostate cancer	41.3%	41.2%	100.0%	41.9%	41.8%	100.0%	38.6%	38.6%	0.0%	38.7%	46.6%	72.2%	53.8%	50.0%
Breast cancer	72.1%	77.8%	72.0%	70.5%	77.8%	70.4%	78.4%	0.0%	78.4%	68.7%	84.6%	73.9%	88.2%	91.7%
Colorectal cancer	62.6%	67.1%	57.9%	62.4%	65.2%	59.4%	62.2%	70.4%	54.1%	60.5%	68.9%	79.5%	69.6%	50.0%
Malignant melanoma of skin	63.5%	63.4%	63.6%	60.7%	58.8%	63.6%	66.7%	71.4%	57.9%	63.6%	14.3%	100.0%	0.0%	0.0%
Cerebrovascular diseases	38.5%	49.3%	30.4%	37.5%	49.0%	29.4%	38.8%	49.1%	30.6%	33.6%	48.8%	64.4%	65.1%	55.0%
Chronic lower respiratory diseases	49.5%	52.0%	47.2%	47.9%	50.5%	45.6%	54.3%	55.9%	52.6%	48.7%	53.8%	70.7%	40.7%	50.0%
Accidents (unintentional injuries)	75.4%	82.7%	63.2%	73.0%	81.3%	60.0%	77.6%	85.2%	65.3%	68.5%	86.2%	86.2%	92.3%	76.2%
Motor vehicle accidents	93.6%	94.8%	91.0%	93.1%	94.2%	90.4%	94.5%	97.0%	89.2%	90.4%	98.0%	100.0%	95.8%	90.0%
Falls	29.5%	41.0%	19.4%	29.4%	41.3%	19.0%	25.8%	37.7%	15.9%	26.4%	40.2%	27.3%	60.0%	42.9%
Accidental poisoning	98.7%	98.9%	98.4%	99.1%	98.9%	99.3%	97.6%	99.2%	94.8%	98.4%	99.4%	100.0%	100.0%	100.0%
Accidental drowning	94.9%	98.2%	87.0%	93.4%	97.6%	84.2%	100.0%	100.0%	100.0%	90.2%	100.0%	100.0%	100.0%	100.0%
Influenza and pneumonia	34.4%	40.1%	28.8%	34.7%	40.0%	29.5%	34.4%	41.3%	27.8%	31.2%	45.7%	40.0%	54.3%	25.0%
Alzheimer's disease	14.5%	19.2%	12.2%	13.9%	18.1%	11.9%	19.7%	27.7%	15.5%	14.3%	14.3%	19.1%	24.0%	22.2%
Diabetes	67.2%	71.6%	61.3%	66.9%	70.8%	62.0%	68.6%	73.5%	60.6%	62.8%	71.8%	74.8%	83.1%	62.5%
Intentional self-harm (suicide)	92.6%	91.6%	96.2%	92.1%	91.4%	94.8%	94.0%	92.1%	100.0%	90.4%	99.3%	100.0%	100.0%	100.0%
Chronic liver disease and cirrhosis	91.6%	94.2%	87.5%	90.5%	93.6%	85.6%	95.2%	97.6%	91.4%	89.4%	91.3%	93.3%	100.0%	100.0%
Nephritis, nephrotic syndrome and nephrosis	43.8%	47.8%	39.7%	39.7%	41.8%	38.0%	48.7%	53.6%	42.6%	36.4%	69.8%	58.8%	67.7%	16.7%
Septicemia	55.6%	55.7%	55.6%	56.5%	54.5%	58.6%	54.9%	63.0%	46.7%	52.3%	61.4%	33.3%	83.3%	100.0%
Assault (homicide)	96.5%	96.9%	95.2%	97.4%	98.4%	93.8%	100.0%	100.0%	100.0%	92.3%	100.0%	100.0%	100.0%	100.0%
Parkinson's disease	28.3%	29.0%	27.0%	28.3%	29.4%	26.4%	27.9%	27.3%	29.4%	26.3%	35.5%	60.0%	33.3%	50.0%
Essential (primary) hypertension and hypertensive renal disease	42.0%	50.3%	34.7%	40.5%	48.2%	33.8%	49.0%	59.2%	38.8%	37.0%	49.0%	67.6%	68.0%	52.4%
Human immunodeficiency virus (HIV) disease	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	0.0%
INJURY BY FIREARMS	91.9%	91.5%	94.0%	91.6%	91.4%	92.4%	94.3%	93.2%	100.0%	88.5%	99.4%	100.0%	100.0%	100.0%
DRUG-INDUCED DEATHS	97.2%	97.4%	96.9%	97.4%	97.6%	97.1%	95.7%	95.8%	95.7%	96.4%	99.4%	100.0%	100.0%	100.0%
ALCOHOL-INDUCED DEATHS	96.1%	95.7%	96.9%	95.5%	95.2%	96.2%	97.9%	97.6%	98.8%	94.5%	96.6%	100.0%	100.0%	100.0%

Notes: ^a Expected years of life at birth for all U.S. residents (78.7 years in 2011); ^b Urban = Maricopa, Pima, Pinal, and Yuma counties.