

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS BY AGE AND GENDER,
ARIZONA, 2013**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	0	*	0	0†	*	*	0	0†	0	0	0	0
5-9	0	*	0	0†	0	0	0	0	0	0	0	0
10-14	9	21	0	30	27	135	0	162	0	0	0	0
15-19	414	702	0	1,116	1,524	6,213	0	7,737	15	6	0	21
20-24	1,059	1,088	0	2,147	3,269	8,960	*	12,231	89	13	0	102
25-29	632	608	0	1,240	1,777	3,677	*	5,456	78	22	0	100
30-34	400	347	0	747	990	1,723	0	2,713	70	8	0	78
35-39	264	183	0	447	444	745	0	1,189	30	*	0	30†
40-44	207	90	0	297	287	403	0	690	46	*	0	50†
45-49	152	57	0	209	194	191	0	385	47	*	0	50†
50-54	122	23	0	145	100	105	0	205	32	*	0	30†
55-59	51	15	0	66	46	42	0	88	18	*	0	20†
60-64	34	*	0	40†	22	12	0	34	7	0	0	7
65-over	14	*	0	20†	17	13	0	30	*	0	0	0†
Total	3,358	3,150†	0	6,510†	8,700†	22,220†	0†	30,920†	440†	60†	0	500†

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.