

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2015**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total					
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate
White Non-Hispanic	386	40.5	10.0	119	22.5	3.1	0	0.0	0.0	1,973	23.9	50.9	5,734	17.6	147.9
Black or African American	117	12.3	36.6	66	12.5	20.6	0	0.0	0.0	1,336	16.2	417.7	2,557	7.9	799.4
Hispanic or Latino	350	36.8	16.7	243	45.8	11.6	0	0.0	0.0	2,431	29.4	115.8	9,901	30.5	471.8
Asian or Pacific Islander	16	1.7	6.5	14	2.6	5.7	0	0.0	0.0	58	0.7	23.7	325	1.0	132.9
American Indian or Alaska Native	42	4.4	14.6	30	5.7	10.4	0	0.0	0.0	526	6.4	183.1	2,430	7.5	845.9
Not Specified	41	4.3	N/A	58	10.9	N/A	0	0.0	N/A	1,946	23.5	N/A	11,555	35.5	N/A
Total	952	100.0	13.9	530	100.0	7.8	0	0.0	0.0	8,270	100.0	121.1	32,511	100.0	476.1

Notes: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2015; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.