

**TABLE 3B-5  
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,  
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2016**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total			Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate			
<b>White Non-Hispanic</b>	452	37.4	11.7	169	24.9	4.4	0	0.0	0.0	2,434	0.2	62.8	6,282	0.2	162.0
<b>Black or African American</b>	139	11.5	43.5	89	13.1	27.8	0	0.0	0.0	1,645	0.2	514.3	2,829	0.1	884.5
<b>Hispanic or Latino</b>	468	38.7	22.3	295	43.5	14.1	0	0.0	0.0	2,931	0.3	139.7	10,118	0.3	482.2
<b>Asian or Pacific Islander</b>	17	1.4	5.9	12	1.8	4.2	0	0.0	0.0	82	0.0	33.5	343	0.0	133.8
<b>American Indian or Alaska Native</b>	98	8.1	40.1	44	6.5	18.0	0	0.0	0.0	658	0.1	229.0	2,432	0.1	846.5
<b>Not Specified</b>	35	2.9	N/A	71	10.3	N/A	0	0.0	N/A	2,580	0.2	N/A	12,935	0.4	N/A
<b>Total</b>	1,209	100.0	17.7	679	100.0	9.9	0	0.0	0.0	10,330	1.0	151.3	32,923	1.0	511.5

Notes: a Number of cases per 100,000 population; Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2016; rates per 100,000 population.

**Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.**