

**TABLE 5B-21
DISTRIBUTION OF LOW-BIRTHWEIGHT (LBW) BIRTHS AND LBW RISK BY
NUMBER OF PRENATAL VISITS AND COUNTY OF RESIDENCE, ARIZONA, 2016**

| | | | Prenatal visits | | | | | | Total |
|------------|---------------------------|----------------------------------|-----------------|------------|------------|-------------|------------|---------|--------|
| | | | No visits | 1-4 visits | 5-8 visits | 9-12 visits | 13+ visits | Unknown | |
| Apache | Births <2,500 grams (LBW) | Count | 6 | 17 | 18 | 25 | 9 | * | 79 |
| | | Share of LBW births ^a | 7.6% | 21.5% | 22.8% | 31.6% | 11.4% | 5.1% | 100.0% |
| | | LBW risk ^b | 31.6% | 24.3% | 7.4% | 5.4% | 4.2% | 30.8% | 7.7% |
| | Total births | Count | 19 | 70 | 242 | 466 | 212 | 13 | 1,022 |
| Cochise | Births <2,500 grams (LBW) | Count | 16 | 23 | 38 | 33 | 17 | * | 130 |
| | | Share of LBW births ^a | 12.3% | 17.7% | 29.2% | 25.4% | 13.1% | 2.3% | 100.0% |
| | | LBW risk ^b | 17.8% | 16.8% | 9.6% | 5.8% | 5.2% | 30.0% | 8.5% |
| | Total births | Count | 90 | 137 | 397 | 572 | 325 | 10 | 1,531 |
| Coconino | Births <2,500 grams (LBW) | Count | * | 17 | 27 | 48 | 31 | 0 | 130† |
| | | Share of LBW births ^a | 3.9% | 13.3% | 21.1% | 37.5% | 24.2% | 0.0% | 100.0% |
| | | LBW risk ^b | 20.0% | 20.0% | 11.1% | 6.6% | 5.9% | 0.0% | 7.9% |
| | Total births | Count | 25 | 85 | 244 | 725 | 528 | 8 | 1,615 |
| Gila | Births <2,500 grams (LBW) | Count | * | 11 | 11 | 15 | 7 | * | 50† |
| | | Share of LBW births ^a | 6.1% | 22.4% | 22.4% | 30.6% | 14.3% | 4.1% | 100.0% |
| | | LBW risk ^b | 10.0% | 13.8% | 9.2% | 6.8% | 5.2% | 18.2% | 8.3% |
| | Total births | Count | 30 | 80 | 119 | 219 | 134 | 11 | 593 |
| Graham | Births <2,500 grams (LBW) | Count | * | 8 | 9 | 17 | 10 | 0 | 50† |
| | | Share of LBW births ^a | 8.3% | 16.7% | 18.8% | 35.4% | 20.8% | 0.0% | 100.0% |
| | | LBW risk ^b | 15.4% | 13.3% | 7.6% | 7.6% | 7.8% | 0.0% | 8.6% |
| | Total births | Count | 26 | 60 | 119 | 224 | 129 | 0 | 558 |
| Greenlee | Births <2,500 grams (LBW) | Count | * | 0 | * | * | * | 0 | 10† |
| | | Share of LBW births ^a | 25.0% | 0.0% | 25.0% | 12.5% | 37.5% | 0.0% | 100.0% |
| | | LBW risk ^b | 100.0% | 0.0% | 5.3% | 1.7% | 7.7% | 0.0% | 5.4% |
| | Total births | Count | * | 11 | 38 | 58 | 39 | * | 150† |
| La Paz | Births <2,500 grams (LBW) | Count | 0 | * | * | * | * | * | 10† |
| | | Share of LBW births ^a | 0.0% | 14.3% | 35.7% | 14.3% | 28.6% | 7.1% | 100.0% |
| | | LBW risk ^b | 0.0% | 15.4% | 8.3% | 2.7% | 5.8% | 33.3% | 6.3% |
| | Total births | Count | * | 13 | 60 | 73 | 69 | * | 220† |
| Maricopa | Births <2,500 grams (LBW) | Count | 120 | 309 | 924 | 1,602 | 860 | 150 | 3,965 |
| | | Share of LBW births ^a | 3.0% | 7.8% | 23.3% | 40.4% | 21.7% | 3.8% | 100.0% |
| | | LBW risk ^b | 14.2% | 15.1% | 12.3% | 6.1% | 5.2% | 14.1% | 7.3% |
| | Total births | Count | 845 | 2,048 | 7,493 | 26,132 | 16,442 | 1,061 | 54,021 |
| Mohave | Births <2,500 grams (LBW) | Count | * | 13 | 38 | 39 | 28 | * | 130† |
| | | Share of LBW births ^a | 3.9% | 10.2% | 29.7% | 30.5% | 21.9% | 3.9% | 100.0% |
| | | LBW risk ^b | 20.8% | 14.9% | 11.7% | 4.9% | 4.9% | 45.5% | 7.1% |
| | Total births | Count | 24 | 87 | 325 | 789 | 567 | 11 | 1,803 |
| Navajo | Births <2,500 grams (LBW) | Count | 15 | 17 | 34 | 45 | 18 | * | 130 |
| | | Share of LBW births ^a | 11.5% | 13.1% | 26.2% | 34.6% | 13.8% | 0.8% | 100.0% |
| | | LBW risk ^b | 45.5% | 14.9% | 10.5% | 6.7% | 5.2% | 10.0% | 8.7% |
| | Total births | Count | 33 | 114 | 325 | 667 | 349 | 10 | 1,498 |
| Pima | Births <2,500 grams (LBW) | Count | 93 | 97 | 268 | 235 | 108 | * | 803 |
| | | Share of LBW births ^a | 11.6% | 12.1% | 33.4% | 29.3% | 13.4% | 0.2% | 100.0% |
| | | LBW risk ^b | 15.7% | 13.6% | 12.2% | 4.9% | 3.6% | 6.3% | 7.0% |
| | Total births | Count | 594 | 713 | 2,192 | 4,843 | 3,029 | 32 | 11,403 |
| Pinal | Births <2,500 grams (LBW) | Count | 6 | 19 | 78 | 103 | 77 | 9 | 292 |
| | | Share of LBW births ^a | 2.1% | 6.5% | 26.7% | 35.3% | 26.4% | 3.1% | 100.0% |
| | | LBW risk ^b | 9.5% | 13.1% | 15.3% | 4.9% | 4.8% | 15.3% | 6.5% |
| | Total births | Count | 63 | 145 | 511 | 2,103 | 1,590 | 59 | 4,471 |
| Santa Cruz | Births <2,500 grams (LBW) | Count | 22 | 7 | 11 | 12 | * | * | 60† |
| | | Share of LBW births ^a | 37.9% | 12.1% | 19.0% | 20.7% | 5.2% | 5.2% | 100.0% |
| | | LBW risk ^b | 31.4% | 8.8% | 6.0% | 5.5% | 4.1% | 15.8% | 9.0% |
| | Total births | Count | 70 | 80 | 183 | 217 | 73 | 19 | 642 |
| Yavapai | Births <2,500 grams (LBW) | Count | * | 16 | 33 | 60 | 31 | * | 140† |
| | | Share of LBW births ^a | 1.4% | 11.1% | 22.9% | 41.7% | 21.5% | 1.4% | 100.0% |
| | | LBW risk ^b | 9.1% | 24.6% | 13.3% | 8.8% | 3.7% | 13.3% | 7.7% |
| | Total births | Count | 22 | 65 | 248 | 680 | 838 | 15 | 1,868 |
| Yuma | Births <2,500 grams (LBW) | Count | 9 | 29 | 67 | 66 | 39 | 0 | 210 |
| | | Share of LBW births ^a | 4.3% | 13.8% | 31.9% | 31.4% | 18.6% | 0.0% | 100.0% |
| | | LBW risk ^b | 5.4% | 14.4% | 12.4% | 5.6% | 4.3% | 0.0% | 7.0% |
| | Total births | Count | 167 | 202 | 539 | 1,189 | 904 | * | 3,004 |
| Unknown | Births <2,500 grams (LBW) | Count | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | Share of LBW births ^a | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | | LBW risk ^b | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | Total births | Count | 0 | 0 | 0 | * | * | 0 | 0† |

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; ^a Percent distribution of LBW births by number of prenatal visits; ^b Number of LBW births per 100 births in specified prenatal care group.