

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2017**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	0	*	*	0†	0	*	0	0†	0	0	0	0
5-9	*	*	0	0†	0	6	0	6	0	*	0	0†
10-14	15	39	0	54	32	155	*	188	0	*	0	0†
15-19	825	1,024	*	1,853	2,267	7,016	19	9,302	62	23	0	85
20-24	1,813	1,407	*	3,224	4,298	10,173	37	14,508	174	70	0	244
25-29	1,651	1,147	*	2,802	2,864	4,857	14	7,735	239	79	0	318
30-34	1,068	699	6	1,773	1,485	2,115	10	3,610	200	65	0	265
35-39	695	429	*	1,128	930	1,137	*	2,070	162	35	0	197
40-44	405	227	*	633	461	512	*	976	106	17	0	123
45-49	333	128	*	462	300	274	*	577	98	11	0	109
50-54	225	72	0	297	213	129	0	342	103	11	0	114
55-59	130	32	*	164	132	61	0	193	56	*	0	60†
60-64	57	11	0	68	49	29	0	78	21	*	0	30†
65-over	44	*	0	50†	33	15	0	48	22	*	0	20†
Total	7,260†	5,220†	30†	12,510†	13,064	26,480†	90†	39,640†	1,243	320†	0	1,560†

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.