

**TABLE 3B-4
RATES^a OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2017**

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS		
	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	0.0	**	**	0.0	**	**	0.0	0.0	0.0
5-9	**	**	**	0.0	2.7	1.3	0.0	**	**
10-14	6.4	17.3	11.8	13.7	68.8	40.7	0.0	**	**
15-19	345.1	450.3	396.4	948.2	3085.0	1989.9	25.9	10.1	18.2
20-24	718.8	605.1	664.3	1703.9	4375.2	2985.2	69.0	30.1	50.3
25-29	657.4	491.7	577.6	1140.4	2082.1	1593.9	95.2	33.9	65.6
30-34	462.4	321.0	393.8	643.0	971.2	802.3	86.6	29.8	59.1
35-39	316.6	200.5	259.3	423.7	531.4	476.8	73.8	16.4	45.4
40-44	196.3	110.5	153.5	223.4	249.2	236.3	51.4	8.3	29.9
45-49	158.4	60.6	109.4	142.7	129.7	136.2	46.6	5.2	25.9
50-54	107.4	33.4	69.8	101.6	59.8	80.4	49.1	5.1	26.8
55-59	62.7	14.3	37.5	63.6	27.2	44.7	27.0	**	13.7
60-64	30.4	5.2	17.0	26.2	13.7	19.5	11.2	**	6.3
65-over	8.2	**	4.2	6.1	2.4	4.1	4.1	**	2.0
Total	211.0	149.8	180.2	379.4	759.2	570.5	36.1	9.2	22.6

Notes: ** Cell suppressed due to rate/ratio/percent based on non-zero count less than 6; ^a Number of cases per 100,000 population; table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2017; denominators for unknown or transgender category are not available; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.