

**TABLE 3B-5  
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,  
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2017**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total			Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate			
<b>White Non-Hispanic</b>	497	31.8	12.7	177	21.3	4.5	0	0.0	0.0	2,598	20.8	66.4	6,703	16.9	171.2
<b>Black or African American</b>	144	9.2	43.8	97	11.7	29.5	0	0.0	0.0	1,671	13.4	508.4	3,048	7.7	927.4
<b>Hispanic or Latino</b>	583	37.3	27.2	336	40.5	15.7	0	0.0	0.0	3,460	27.6	161.3	11,733	29.6	547.1
<b>Asian or Pacific Islander</b>	18	1.2	7.2	11	1.3	4.4	0	0.0	0.0	85	0.7	34.0	354	0.9	141.7
<b>American Indian or Alaska Native</b>	141	9.0	48.0	50	6.0	17.0	0	0.0	0.0	845	6.8	287.7	2,354	5.9	801.6
<b>Not Specified</b>	181	11.6	N/A	159	19.2	N/A	0	0.0	N/A	3,855	30.8	N/A	15,443	39.0	N/A
<b>Total</b>	1,564	100.0	22.6	830	100.0	12.0	0	N/A	N/A	12,514	100.0	180.5	39,635	100.0	571.8

Notes: a Number of cases per 100,000 population; Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2017; rates per 100,000 population.

**Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.**