



1D.

ABORTIONS


In July 1976, the Arizona Department of Health Services (ADHS) began an Abortion Surveillance Program (*Arizona Administrative Code, R9-19-302D*, as authorized by the *Arizona Revised Statute 36-344*). The program required the reporting of induced terminations of pregnancy performed in hospitals, outpatient treatment centers, and physicians' offices throughout the State.

In 2010, SB1304 was passed by the legislature that statutorily required the reporting of abortions and treatment of complications associated with abortions. In addition, the office of the Courts were required to report the number of petitions received by the court for authorization to perform abortions on unemancipated minors as prescribed in Arizona Revised Statutes §36-2152. The collection of this information began on July 29, 2010.

Based on the information collected on race/ethnicity in the abortion reporting system, the racial/ethnic groups described in this section differ from that of other sections. In 2017, 4.7 percent (577) of abortion records had race and ethnicity identified as unknown, and 1.8 percent (216) of abortion records had two or more racial/ethnic groups identified. Due to the proportion of information on race/ethnicity that could not be categorized by the five racial/ethnic groups used through this report, this section reports numbers for records with unknown or multiple race information as separate groups.

1D. ABORTIONS

Figure 1D-1
Number of Reported Abortions and Abortion Rates^a by Year,
Arizona Residents, 2007-2017




In calendar year 2017, the Arizona Department of Health Services received 12,336 reports of abortions obtained by Arizona residents, 834 less than the 13,170 reported in 2016 (**Figure 1-D1, Table 1D-1**).

There were 12,301 abortions in non-hospital clinics (**Table 1D-1**). Hospitals accounted for 31 abortion procedures, with less than 6 procedures being performed outside of clinics and hospitals.

Ninety-one percent of all reported abortions occurred at or before 13 weeks of pregnancy. Few reported abortions occurred after 14 or more weeks of gestation: 8.0 percent at 14 weeks or more (percentages based on data in **Table 1D-3**).

Notes: ^a Number of abortions per 1,000 females 15-44 years old. The sharp increase in number of reported abortions and rates was due to enhanced surveillance as noted in the abortion report.

Figure 1D-2
Trends in the Number of Surgical and Non-Surgical Abortions by Year,
Arizona Residents, 2007-2017




The rate of induced terminations of pregnancy per 1,000 female residents aged 15-44 decreased from 10.0 in 2016 to 9.2 in 2017 (**Table 1D-1**).

Trends in the percent of abortions conducted through surgical and non-surgical from 2007 to 2017 indicate shifting practices in methods used to complete abortions. In 2007, 64.1 percent of all abortion procedures were surgical procedures. By 2009, the number of abortions performed by surgical versus non-surgical means was nearly equal. Since then, the proportion of abortions conducted through surgical means has again increased, with 63.3 percent of all abortion procedures conducted in 2017 being surgical procedures.

1D. ABORTIONS

Figure 1D-3
Abortion Ratios^a by Age Group, Arizona Residents, 2017




Abortion ratios (the number of abortions per 1,000 live births in each age group) vary by age at reported termination of pregnancy. Ratios tend to be higher at the extremes of the age distribution of the childbearing period, i.e., among women under age 15 and those aged 40 years or older (**Figure-1D-3**).

A comparison of abortion ratios by race/ethnicity is provided in **Table 1D-2**.

Note: ^a Number of reported abortions per 1,000 live births in specified group.

Figure 1D-4
Proportional Contribution of Non-Surgical^a and Surgical Abortions by Race/Ethnicity, Arizona Residents, 2017



In 2017 non-surgical (also called “medical”) abortions made up 36.7 percent of all procedures reported in Arizona. Among Asians and White non-Hispanics, the proportion of non-surgical abortions exceeded the average for all groups (**Figure 1D-4**). The proportion of non-surgical abortions among Black or African American women was the lowest among racial/ethnic groups.

Notes: ^a Non-surgical: Mifepristone, Methotrexate, RU486, etc.; Reporting changes related to race/ethnicity makes comparisons with previous years difficult.