

3B.

SEXUALLY TRANSMITTED DISEASES


All states require that certain sexually transmitted diseases (STDs) be reported by physicians and other health care providers when they suspect that a case has occurred or they have laboratory confirmation.

It is important to note that disease reporting is likely incomplete and completeness may vary depending on the disease. Moreover, changes in methods for public health surveillance, or implementation of new diagnostic tests can cause changes in disease reporting that are independent of the true incidence of disease.* In this section, rates for STDs were calculated using denominators based on 2016 estimates taken from the CDC.

*Centers for Disease Control and Prevention. Summary of notifiable diseases – United States, 2008. Published June 25, 2010, for 2008; Vol. 57 (No. 54). Available online at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5754a1.htm>

3B. SEXUALLY TRANSMITTED DISEASES

Figure 3B-1
Trends in the Incidence Rates^a of Gonorrhea by Year, Arizona, 2007-2017




Neisseria gonorrhoeae infection is the second most commonly reported notifiable disease in the United States. (Figure 3B-1). The consistent steady increase in the incidence rate of gonorrhea since 2009 likely resulted from a combination of factors, such as changes in surveillance, increases in the number of tests performed, and actual increases in disease occurrence (Figure 3B-1).

The *Healthy People 2010* objective HP25-2 defines the target rate for gonorrhea as equal to or lower than 19.1 cases per 100,000 population. However, the *Healthy People 2020* target is for ages 15-44 and is set at 151.0/100,000 females and 147.0/100,000 males (Table 6A-2).

Note: ^a Number of reported cases per 100,000 population.

Figure 3B-2
Trends in the Incidence Rates^a of Gonorrhea among Females 15-24 and 15-44 Years, Arizona, 2007-2017



The 2017 incidence rate for gonorrhea was 370.8 per 100,000 for Arizona females aged 15-44 years, meaning Arizona's incidence rate was higher than the *Healthy People 2020* objective. Generally, the trend in gonorrhea incidence rates are similar for women aged 15-24 and aged 15-44, although the overall incidence rate is consistently higher for women aged 15-24.


Notes: ^a Number of reported cases per 100,000 females; There was a change in target rate and age range for Healthy People 2020 objective. In National Year 2010 objective was for females ages 15-24. In Healthy People 2020 objective is for females ages 15-44.

3B. SEXUALLY TRANSMITTED DISEASES

Chlamydia trachomatis is the most prevalent bacterial sexually transmitted disease in the United States (1,708,569 cases in 2017) with the highest rates reported among adolescents and young adults (Table 3B-4). Recent availability of sensitive tests for *chlamydia* using DNA amplification technology undoubtedly contributed to the increase in the number of reported cases in Arizona over the last decade (Figure 3B-3, Table 3B-1).

The incident rate of chlamydia was previously reported for females 15-24 years, however based on changes in *Healthy People 2020*, it would be reported for females 15-44 years. The *Healthy People 2020* goal for chlamydia is set at 1,478.2 per 100,000 females. The incidence rate for Arizona in 2017 was 1,940 per 100,000 females age 15-44 years (Table 6A-2).

Figure 3B-3
Trends in the Incidence Rates^a of Chlamydia among Females 15-24 and 15-44 Years, Arizona, 2007-2017



<http://www.cdc.gov/std/stats17/chlamydia.htm>

Notes: ^a Number of reported cases per 100,000 females; There was a change in target rate and age range for Healthy People 2020 objective. In National Year 2010 objective was for females ages 15-24. In Healthy People 2020 objective is for females ages 15-44.

Congenital syphilis is an infection caused by the spirochete *Treponema pallidum*, which can be passed from the mother to child during fetal development or birth. Not all infants born to infected women will be infected.

In 1988, CDC implemented a new Congenital syphilis case definition. It no longer relies on documentation of infection in the infant; rather, it presumes that an infant is infected if it cannot be proven that an infected mother was adequately treated for syphilis before or during pregnancy.

The *Healthy People 2020* goal for congenital syphilis is 14.5 cases per 100,000. In Arizona, the incidence rate of congenital syphilis has seen a sharp increase at 36.7/100,000 in 2017 following a period of consistent decreases (Figure 3B-4, Table 6A-2).

Figure 3B-4
Trends in the Incidence Rates^a of Congenital Syphilis by Year, Arizona, 2007-2017

