

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2018**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total			Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate			
White Non-Hispanic	686	35.2	17.4	322	25.9	8.2	0	0.0	0.0	3,229	25.0	81.8	8,599	21.0	217.8
Black or African American	199	10.2	58.5	141	11.3	41.5	0	0.0	0.0	2,152	16.7	632.6	3,934	9.6	1156.5
Hispanic or Latino	729	37.4	32.7	523	42.0	23.5	0	0.0	0.0	3,408	26.4	153.0	11,822	28.9	530.7
Asian or Pacific Islander	33	1.7	12.6	14	1.1	5.3	0	0.0	0.0	123	1.0	46.9	460	1.1	175.3
American Indian or Alaska Native	234	12.0	78.3	141	11.3	47.2	0	0.0	0.0	1,157	9.0	387.4	3,082	7.5	1031.9
Multi-racial	45	2.3	N/A	21	1.7	N/A	0	0.0	0.0	150	1.2	N/A	236	0.6	N/A
Not Specified	25	1.3	N/A	83	6.7	N/A	0	0.0	N/A	2,684	20.8	N/A	12,733	31.2	N/A
Total	1,951	100.0	27.6	1,245	100.0	17.6	0	N/A	N/A	12,903	100.0	182.3	40,866	100.0	577.5

Notes: ^a Number of cases per 100,000 population; Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2018; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.