

**TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED^a YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY
GENDER, RACE/ETHNICITY, AND URBAN/RURAL AREAS^b, ARIZONA, 2019**

	ARIZONA			Area/gender						Race/ethnicity				
	Total	Male	Female	Total Urban	Urban Male	Urban Female	Total Rural	Rural Male	Rural Female	White Non-Hispanic	Hispanic or Latino	Black or African American	American Indian or Alaska Native	Asian or Pacific Islander
Total, all causes	55.6%	61.9%	48.2%	54.8%	61.3%	47.3%	58.2%	63.4%	51.6%	50.8%	67.3%	75.3%	77.8%	60.2%
Major cardiovascular diseases	44.3%	52.3%	34.8%	43.4%	51.8%	33.9%	47.4%	54.0%	38.7%	40.5%	53.8%	70.8%	63.9%	50.9%
Diseases of heart	44.8%	52.4%	34.7%	43.8%	51.9%	33.6%	48.0%	53.7%	39.5%	40.9%	55.2%	70.8%	66.3%	53.1%
Coronary heart disease	46.2%	53.5%	35.3%	45.2%	53.2%	33.8%	49.5%	54.4%	41.3%	42.5%	57.4%	68.7%	66.7%	57.9%
Malignant neoplasms	66.8%	66.7%	66.9%	66.6%	66.6%	66.7%	67.1%	66.6%	67.7%	64.0%	75.7%	76.9%	81.2%	73.9%
Lung cancer	67.5%	70.2%	64.4%	66.5%	68.8%	63.8%	70.4%	74.6%	66.2%	67.0%	67.1%	76.1%	75.0%	60.4%
Colorectal cancer	67.7%	72.0%	62.3%	69.5%	74.3%	63.6%	61.0%	63.8%	57.3%	63.9%	77.7%	84.6%	81.3%	75.0%
Breast cancer	72.3%	83.3%	72.2%	71.7%	83.3%	71.5%	74.9%	0.0%	74.9%	69.3%	75.6%	81.5%	91.7%	100.0%
Prostate cancer	46.3%	46.3%	0.0%	46.0%	46.0%	0.0%	47.0%	47.0%	0.0%	45.5%	48.8%	58.5%	46.7%	14.3%
Malignant melanoma of skin	64.6%	62.4%	68.5%	64.6%	62.6%	68.4%	65.9%	61.5%	73.3%	62.0%	100.0%	100.0%	100.0%	0.0%
Accidents (unintentional injuries)	79.0%	85.1%	66.6%	78.4%	84.8%	65.2%	80.7%	85.4%	71.7%	71.7%	91.6%	96.0%	93.8%	78.0%
Accidental poisoning	99.8%	99.9%	99.6%	99.9%	100.0%	99.7%	99.2%	99.4%	98.8%	99.8%	100.0%	100.0%	99.2%	100.0%
Falls	32.3%	40.9%	23.4%	30.2%	39.0%	21.1%	41.2%	49.5%	33.0%	28.9%	46.8%	62.5%	63.9%	16.7%
Motor vehicle accidents	92.4%	93.1%	90.6%	91.7%	92.4%	90.0%	93.9%	94.8%	92.0%	88.6%	96.6%	100.0%	97.2%	94.1%
Accidental drowning	92.0%	98.6%	75.9%	89.2%	98.0%	69.6%	100.0%	100.0%	100.0%	86.9%	100.0%	100.0%	100.0%	100.0%
Chronic lower respiratory diseases	50.5%	53.1%	48.1%	49.5%	52.3%	47.0%	53.0%	54.7%	51.2%	49.6%	46.6%	69.5%	67.5%	48.0%
Alzheimer's disease	16.4%	20.7%	14.1%	16.7%	22.1%	13.9%	14.7%	13.9%	15.1%	16.3%	16.8%	13.8%	15.8%	17.1%
Cerebrovascular diseases	40.4%	48.7%	34.0%	39.6%	47.2%	33.8%	43.4%	53.4%	34.7%	36.3%	49.5%	73.0%	61.7%	44.0%
Diabetes	71.9%	75.0%	67.3%	72.1%	74.1%	69.2%	70.8%	78.3%	58.8%	68.9%	74.2%	78.2%	79.8%	73.3%
Intentional self-harm (suicide)	90.9%	89.7%	94.9%	91.5%	90.5%	94.7%	88.8%	87.0%	95.5%	88.7%	96.7%	97.8%	98.8%	90.5%
Chronic liver disease and cirrhosis	91.8%	92.2%	91.2%	91.6%	91.7%	91.2%	92.3%	93.2%	91.2%	90.5%	90.4%	100.0%	96.8%	80.0%
Essential (primary) hypertension and hypertensive renal disease	44.0%	54.0%	35.0%	43.6%	54.3%	34.0%	4.5%	51.3%	40.5%	40.7%	49.1%	64.9%	53.2%	37.0%
Influenza and pneumonia	52.0%	54.0%	49.5%	50.1%	52.5%	47.4%	57.3%	58.3%	56.0%	48.6%	57.4%	83.3%	60.7%	45.5%
Parkinson's disease	29.7%	30.9%	27.2%	29.0%	29.9%	27.4%	31.9%	35.2%	26.4%	29.2%	31.3%	0.0%	34.8%	33.3%
Nephritis, nephrotic syndrome and nephrosis	53.6%	57.0%	49.1%	52.1%	54.2%	49.4%	58.4%	64.9%	46.9%	48.2%	64.2%	74.1%	62.3%	52.9%
Septicemia	69.9%	66.7%	73.6%	71.7%	68.8%	74.8%	64.8%	62.3%	68.4%	64.4%	78.3%	80.0%	86.1%	66.7%
Assault (homicide)	98.0%	98.4%	96.7%	98.0%	98.2%	97.3%	98.2%	100.0%	94.7%	94.4%	100.0%	100.0%	100.0%	100.0%
Human immunodeficiency virus (HIV) disease	97.1%	96.2%	100.0%	96.4%	95.3%	100.0%	100.0%	100.0%	100.0%	94.7%	100.0%	100.0%	100.0%	0.0%
Injury by firearms	91.3%	90.3%	96.1%	92.3%	91.5%	96.5%	87.1%	85.6%	94.6%	87.8%	98.3%	98.6%	100.0%	100.0%
Drug-induced deaths	98.7%	99.1%	97.7%	99.0%	99.5%	97.6%	97.0%	96.3%	98.1%	98.1%	99.8%	99.2%	99.0%	100.0%
Opioid-induced deaths	99.3%	99.5%	98.9%	99.5%	99.8%	98.7%	97.8%	96.5%	100.0%	98.9%	100.0%	100.0%	100.0%	100.0%
Alcohol-induced deaths	96.5%	95.8%	98.3%	96.7%	95.7%	98.9%	96.2%	95.8%	97.0%	96.3%	94.1%	100.0%	98.9%	75.0%

Notes: ^a Expected years of life at birth for all U.S. residents (78.8 years in 2019); ^b Urban = Maricopa, Pima, Pinal, and Yuma counties.