

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2019**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	*	*	0	10†	*	*	0	10†	0	0	0	0
5-9	*	*	0	10†	0	*	0	0†	0	0	0	0
10-14	15	24	0	39	37	144	0	181	*	0	0	0†
15-19	785	1,040	*	1,826	2,402	7,242	*	9,646	62	42	0	104
20-24	2,109	1,714	*	3,826	5,119	10,701	10	15,830	296	108	0	404
25-29	2,022	1,412	0	3,434	3,317	5,067	*	8,386	398	122	0	520
30-34	1,456	902	*	2,361	1,818	2,331	7	4,156	321	110	0	431
35-39	971	541	0	1,512	1,120	1,140	*	2,263	251	68	0	319
40-44	531	303	*	835	604	618	0	1,222	168	45	0	213
45-49	415	163	0	578	389	305	0	694	144	24	0	168
50-54	319	75	*	395	258	152	0	410	117	12	0	129
55-59	206	49	0	255	162	74	0	236	100	9	0	109
60-64	88	18	0	106	76	33	0	109	45	*	0	50†
65-over	61	10	0	71	58	18	0	76	37	*	0	40†
Total	8,980†	6,260†	10†	15,250†	15,360†	27,830†	20†	43,220†	1,940†	550†	0	2,490†

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.