

**TABLE 3B-5  
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,  
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2019**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total					
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate
<b>White Non-Hispanic</b>	755	30.4	19.0	338	23.4	8.5	0	0.0	0.0	3,360	22.0	84.4	7,797	18.0	195.9
<b>Black or African American</b>	276	11.1	78.4	144	10.0	40.9	0	0.0	0.0	2,439	16.0	692.7	4,085	9.5	1160.1
<b>Hispanic or Latino</b>	992	39.9	43.5	606	41.9	26.6	0	0.0	0.0	4,258	27.9	186.8	12,507	28.9	548.7
<b>Asian or Pacific Islander</b>	39	1.6	14.1	16	1.1	5.8	0	0.0	0.0	137	0.9	49.4	504	1.2	181.6
<b>American Indian or Alaska Native</b>	299	12.0	100.0	181	12.5	60.5	0	0.0	0.0	1,076	7.1	359.7	2,353	5.4	786.6
<b>Multi-racial</b>	31	1.2	N/A	13	0.9	N/A	0	0.0	0.0	110	0.7	N/A	180	0.4	N/A
<b>Not Specified</b>	93	3.7	N/A	148	10.2	N/A	0	0.0	N/A	3,869	25.4	N/A	15,794	36.5	N/A
<b>Total</b>	2,485	100.0	34.6	1,446	100.0	20.1	0	N/A	N/A	15,249	100.0	212.1	43,220	100.0	601.2

Notes: <sup>a</sup> Number of cases per 100,000 population; table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2019; rates per 100,000 population.

**Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.**