

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2020**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	*	6	0	10†	*	7	0	10†	0	0	0	0
5-9	*	*	0	10†	0	*	0	0†	0	0	0	0
10-14	11	42	0	53	26	156	0	182	0	0	0	0
15-19	885	1,202	*	2,090	1,974	6,170	8	8,152	60	44	0	104
20-24	2,121	1,845	*	3,970	4,304	9,280	*	13,589	289	133	0	422
25-29	2,117	1,318	*	3,438	2,786	4,382	*	7,173	389	129	0	518
30-34	1,582	950	*	2,535	1,616	2,068	*	3,686	365	120	0	485
35-39	1,032	675	0	1,707	947	1,039	*	1,989	285	81	0	366
40-44	599	336	*	936	484	440	*	925	190	37	0	227
45-49	421	190	0	611	331	217	0	548	144	26	0	170
50-54	283	101	*	385	220	118	*	339	141	22	0	163
55-59	206	52	0	258	140	75	0	215	118	8	0	126
60-64	99	14	0	113	70	27	0	97	59	*	0	60†
65-over	60	10	0	70	51	18	0	69	36	0	0	36
Total	9,420†	6,740†	20†	16,180†	12,950†	24,000†	30†	36,980†	2,076	600†	0	2,679

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.