TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2020

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total			CHLAMYDIA		
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate
White Non-Hispanic	820	30.6	20.7	335	20.5	8.4	0	0.0	0.0	3,227	19.9	81.4	6,037	16.3	152.3
Black or African American	317	11.8	87.7	175	10.7	48.4	0	0.0	0.0	2,743	17.0	758.5	3,569	9.7	986.9
Hispanic or Latino	1,007	37.6	44.2	639	39.2	28.1	0	0.0	0.0	4,409	27.2	193.6	9,840	26.6	432.0
Asian or Pacific Islander	35	1.3	12.4	20	1.2	7.1	0	0.0	0.0	116	0.7	41.2	332	0.9	118.0
American Indian or Alaska Native	340	12.7	116.9	289	17.7	99.4	0	0.0	0.0	1,163	7.2	399.8	2,104	5.7	723.3
Multi-racial	45	1.7	N/A	37	2.3	N/A	0	0.0	0.0	265	1.6	N/A	382	1.0	N/A
Not Specified	115	4.3	N/A	136	8.3	N/A	0	0.0	N/A	4,257	26.3	N/A	14,713	39.8	N/A
Total	2,679	100.0	37.3	1,631	100.0	22.7	0	N/A	N/A	16,180	100.0	225.5	36,977	100.0	515.3

Notes: ^a Number of cases per 100,000 population; table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2020; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.