

**TABLE 8-10
SELECTED SUBSTANCES INVOLVED IN DRUG POISONING DEATHS, ARIZONA, 2014**

| | Total | Gender | | Race/ethnicity | | | | | | Age group | | | | | | Marital status | | | | |
|---|-------|--------|-----|--------------------|--------------------|---------------------------|----------------------------------|---------------------------|----|-----------|-------|-------|-------|-----|---------------|----------------|---------|----------|---------|--|
| | | M | F | White non-Hispanic | Hispanic or Latino | Black or African American | American Indian or Alaska Native | Asian or Pacific Islander | <1 | 1-14 | 15-19 | 20-44 | 45-64 | 65+ | Never married | Married | Widowed | Divorced | Unknown | |
| Heroin | 180† | 151 | 29 | 126 | 39 | 6 | 8 | * | 0 | 0 | * | 122 | 52 | * | 128 | 20 | * | 24 | 6 | |
| Other opioids (e.g., codeine, morphine, oxycodone) | 290† | 143 | 147 | 226 | 45 | 10 | 8 | * | 0 | * | * | 102 | 146 | 37 | 91 | 78 | 20 | 96 | * | |
| Methadone | 60† | 35 | 28 | 48 | 13 | * | 0 | 0 | 0 | 0 | * | 23 | 36 | * | 25 | 15 | * | 19 | 0 | |
| Other synthetic narcotics | 50† | 27 | 27 | 39 | 9 | * | * | 0 | 0 | * | 0 | 22 | 23 | 8 | 17 | 16 | * | 17 | * | |
| Cocaine | 50† | 32 | 20 | 30 | 11 | 8 | * | * | 0 | 0 | 0 | 20 | 28 | * | 20 | 16 | * | 13 | * | |
| Other and unspecified narcotics | 30† | 23 | 11 | 28 | * | * | 0 | 0 | 0 | 0 | * | 13 | 15 | * | 15 | 10 | 0 | 9 | 0 | |
| Tricyclic and tetracyclic antidepressants | 40† | 17 | 21 | 28 | * | * | * | 0 | 0 | 0 | * | 14 | 19 | * | 7 | 16 | * | 12 | 0 | |
| Other/unspecified antidepressants | 80† | 25 | 58 | 64 | 10 | * | * | 0 | 0 | 0 | * | 31 | 39 | 12 | 21 | 24 | * | 34 | * | |
| Psychostimulants (e.g., methamphetamine) | 250† | 186 | 63 | 181 | 43 | 9 | 15 | * | 0 | 0 | * | 118 | 125 | * | 119 | 39 | 6 | 66 | 19 | |
| Other/unspecified psychotropic drugs | 0† | 0 | * | * | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | * | 0 | 0 | 0 | * | 0 | |
| Barbiturates | 10† | * | 6 | 8 | * | 0 | 0 | 0 | 0 | 0 | * | * | * | * | * | * | * | * | 0 | |
| Benzodiazepines | 120† | 61 | 61 | 100 | 14 | * | * | * | 0 | * | * | 48 | 61 | 11 | 43 | 37 | 8 | 32 | * | |
| Other antiepileptic and sedative-hypnotic drugs | 20† | 11 | 9 | 12 | 7 | 0 | * | 0 | 0 | 0 | 0 | 9 | 8 | * | * | 8 | * | 7 | 0 | |
| Other and unspecified | 780† | 407 | 371 | 619 | 86 | 30 | 36 | 7 | 0 | * | 7 | 328 | 355 | 86 | 280 | 198 | 55 | 221 | 24 | |

Notes: * Cell suppressed due to count less than 6; † Sum rounded to nearest tens unit due to addend less than 6; A poisoning death can have one or multiple substances listed on the death certificate; The substances in the above table were identified using the ICD-10 codes T36.0 - T50.9; However, these codes are not used for coding of the underlying cause of death; Rather, the corresponding external cause code will be designated as the underlying cause (see Table 8-1).